

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

"Made available under NASA sponsorship
in the interest of early and wide dis-
semination of Earth Resources Survey
Program information and without liability
for any use made thereof."

E7.6-10333

CR-147214

A STUDY OF THE HAOR AREAS OF SYLHET-MYMENSING
DISTRICTS WITH ERTS IMAGERIES (WINTER CROP ESTIMATION)

RECEIVED
NASA STI FACILITY
ACQ. BR.

JUN 08 1976

DCAF# 1612911010

by

A. M. Choudhury, A. Azim,
S. Ahmed and S. Rahman,
Bangladesh ERTS Programme.

Pi - ANWAR HUSSAIN

(E76-10333) A STUDY OF THE HAOR AREAS OF
SYLHET-MYMENSING DISTRICTS WITH ERTS
IMAGERIES (WINTER CROP ESTIMATION) (Atomic
Energy Commission) 5 p HC \$3.50 CSCL 02C

N76-23659

G3/43 00333

Unclas

Presented at the First Annual Bangladesh Science Conference,.

March 28 - March 31, Dacca 1976.

A STUDY OF THE HAOR AREAS OF SYLHET-MYMENSING
DISTRICTS WITH ERTS IMAGERIES(WINTER-CROP ESTIMATION)

The advent of Space Age has opened a new era in the exploration of the earth. In the beginning the emphasis was more on the studying the heaven from above the earth's atmosphere. But it has been found that we can study the earth also better from heaven. Since the 1st Sputnik was launched on 4th October, 1957, more than 1400 spacecrafts have been thrown into space. Of all the direct application satellite systems envisaged, those for measuring the resources of the earth show by far the greatest promise of benefit to the world.

ERTS Programme

The meteorological satellites became operational in early sixties. These satellites and the manned space flight programmes namely Mercury, Gemini and Apollo demonstrated that not only the weather of the earth but various features of the earth can be detected by means of suitable sensors placed on board the satellite. To realise this end, NASA launched the first ERTS (Earth Resources Technology Satellite) later named Landsat on July 23, 1972. Landsat-2 was launched on 22 January, 1975. The method is often called Remote Sensing as the information which is obtained about an object or phenomenon is not in direct contact with the information gathering device. Both these satellites have got seven channels or bands as they are called. The frequency ranges are as follows:-

RBV (Return Beam Vidicon)

	Wave length in micrometer
Band 1	0.475-.575
Band 2	0.580-.680
Band 3	0.690-.830

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR

MSS (Multispectral Scanner)

Band 4	0.5-0.6
Band 5	0.6-0.7
Band 6	0.7-0.8
	0.8-1.1
Band 7	

... up a plan of participating in the ERTS Programme Bangladesh the following sectors (1) Agriculture (2) Water Resources (3) Forestry (4) Fisheries and Oceanography (5) Geology (6) Cartography (7) Meteorology (8) Interpretation Techniques Development. Presently the Programme is sponsored by the Planning Commission.

Data Interpretation

All objects like water, vegetation, soil etc radiate electromagnetic energy and receive radiation from their surroundings. The electromagnetic energy radiated depends on the area, nature and temperature of the radiating surface. This can be detected by means of suitable sensors on board the satellite and is called the spectral signature of the body. This property of physical objects forms the basis of multispectral data analysis in Remote Sensing. (SLIDES)

Spectral response of a particular object is not unique. Thus there may be an overlap in the spectral signature of two bodies. The measurement vector corresponding to a particular ground resolution element from one part of a field will not correspond exactly to the measurement vector corresponding to another ground resolution element from the same field. Vectors from the same class tend to form a cluster of points. In the multispectral approach, if there are n channels, we get n number of measurements for a particular ground resolution element.

They define a point in n dimensional Euclidean Space called Measurement Space, represented by a vector $X=(x_1, x_2, \dots, x_n)$. This n dimensional measurement vector is converted into an n dimensional feature vector $Y=(y_1, y_2, \dots, y_n)$. This feature vector is then assigned a class based on Statistical Decision Theory. Each point in the feature space is assigned a class based on some criterion depending on the spectral characteristics.

Analysis of the Sylhet-Mymensingh Haor Frame

The satellite takes picture from a height of 560 miles. Each frame covers a distance of 115×115 miles. Same spot is repeated after every eighteen days. The slides show the SYLHET-MYMENSINGH Haor area in Bands 5 and 7.

We have chosen this area as an experimental site for winter crop estimation. We have visited a few test sites just to find out what tone represents what in the actual field in different bands. This process is called finding the ground truth. We have selected band 7 and band 5 for 27/2 March, 1975. The boro crop was identified in both the bands. A very rough estimate gives an acreage of 11.5 acres. The Agriculture Department Figure for 1973 is 12.3 acres. Our estimate is about 93% of the 1973 figure which is available to us. Perhaps we have not taken into account very small fields. Plots less than five acres cannot be identified in ERTS imageries. This shows the feasibility of objective crop estimation with the help of satellites. This has been done roughly with the unaided eye. With the help of computers, it is possible to get an accuracy of 98% or more.

REPRODUCIBILITY OF THE
ORIGINAL PAGE IS POOR


Jahangirnagar University and the ERTS group have undertaken a joint programme with the Environmental Research Institute of Michigan (ERIM) for mapping this region with the help of their computer and when we get their results, we shall be able to compare the results. We will also visit the test areas again in April just to check our results.

Haor area is a very resourceful area but very undeveloped. A study of the area with the satellite data will be very helpful for the optimal utilization of this area.

Acknowledgement: The authors are grateful to Dr. Anwar Hossain, Principal Investigator, Bangladesh ERTS Programme and Prof. M.I. Chowdhury, Chairman, Geography Department, Jahangirnagar University for encouragement and valuable discussions.

References:

1. Wilfred E. Scull: The Earth Resources Technology Satellites ERTS A and B, NASA, 1972.
2. Philip H. Swain: Pattern Recognition: A Basis for Remote Sensing Data Analysis, LARS Information Note 11572.
3. A.M. Choudhury: Introduction to Mathematical Approach to ERTS Data Interpretation: Proceedings of the Remote Sensing Seminar, Dacca, 1975.
4. Dr. Anwar Hossain: Bangladesh National Programme of the Earth Resources Technology Satellite ERTS-B, 1974.
5. Bangladesh Agriculture in Statistics: Statistical Series No.1, Ministry of Agriculture, Govt. of the People's Republic of Bangladesh, 1973.



ORIGINAL PAGE IS
OF POOR QUALITY

